



[Home](#)

[Search](#)

[Causes of prostatitis](#)

[Drainage in Prostatitis](#)

[Methods of treatment](#)

[Alpha List of files](#)

[Welcome doctors](#)

[Prostatitis Foundation,](#)

[Pointers to other sites](#)

[Archive](#)

# Prostatitis

Antimicrobial Agents and Chemotherapy, February 2000, p. 261-266, Vol. 44, No. 2

## Ultrastructure of Achilles Tendons of Rats Treated with Ofloxacin and Fed a Normal or Magnesium-Deficient Diet

Mehdi Shakibaei, <sup>1</sup>Kerstin Pfister, <sup>2</sup>Rudolf Schwabe, <sup>2</sup>Jürgen Vormann,<sup>3</sup> and Ralf Stahlmann <sup>2,\*</sup>, <sup>1</sup>Institute of Anatomy, <sup>1</sup>Institute of Clinical Pharmacology and Toxicology, <sup>2</sup>and Institute of Molecular Biology and Biochemistry, <sup>3</sup>Benjamin Franklin Medical Center, Freie Universität Berlin, 14195 Berlin, Germany

### ABSTRACT

Fluoroquinolones can cause tendinitis and tendon rupture. However, toxicological as well as clinical information on quinolone-induced tendopathy is scarce. We performed extensive electron microscopic studies with Achilles tendon specimens from ofloxacin-treated rats. The drug was given at a dose of 1,200 mg/kg (body weight) orally. Juvenile Wistar rats received one or three oral doses each of 1,200 mg of ofloxacin/kg (body weight)/day.

Three days after treatment, the tenocytes of their Achilles tendons showed degenerative alterations, such as multiple vacuoles and vesicles in the cytoplasm that had developed due to swellings and dilatations of cell organelles. Other indications of cell degradation were the occurrence of cell debris and cell detachment from the extracellular matrix accompanied by a loss of cell-matrix interaction.

The tenocytes of juvenile Wistar rats that had been treated at day 36 with a single oral dose of 1,200 mg of ofloxacin/kg (body weight) and sacrificed either 3 or 6 months later exhibited similar degenerative alterations. The number of degenerative alterations of tenocytes after ofloxacin treatment was considerably higher in rats that had received a magnesium-deficient diet than in rats with normal magnesium status. Of the adult rats that had been treated once, 5 times, and 10 times with ofloxacin and killed 1 day later, only those with the 10-times treatment showed a significantly increased number of degeneratively altered tenocytes. In summary, effects observed in tendons show similar pathological features as described earlier in cartilage, indicating that quinolone-induced arthropathy and quinolone-induced tendopathy probably are different clinical manifestations of the same toxic effect on cellular components of connective tissue structures.

The full article is copyrighted by the publisher. If you to to the [American Microbiological Society website](#) , you can find out how to obtain the full article.

---

This information is forwarded to you by the Prostatitis Foundation. We do not provide medical advice. We distribute literature and information relevant to prostatitis. While we encourage all research we do not endorse any doctor, medicine or treatment protocol. Consult with your own physician.

© 2002 The Prostatitis Foundation



[Further Contact:](#)(click on words or mailbox)

This page was created by [Ideasmith®](#).

[Add to this site](#)